

Hepatitis C

Quick reference guide for primary care

What is hepatitis C (HCV)?

- A blood-borne virus, spread mainly through blood-to-blood contact
- Can damage the liver, potentially causing cirrhosis and primary liver cancer
- Symptoms can take years or decades to occur
- An estimated 200,000 people are chronically infected in England
- HCV has been associated with injecting drug use, but there are a variety of ways in which it can be transmitted (see below)

Why should I be proactive in diagnosing HCV?

- About half of those infected in England are probably unaware of it
- Treatment can successfully clear the virus in more than half of patients treated overall

Who is at risk of HCV?

Hepatitis C testing should also be offered to anyone who:

- Has unexplained abnormal liver function tests (e.g. elevated ALT), or unexplained jaundice
- Has ever injected drugs in the past (including anabolic steroids) using shared equipment, however long ago, even if this was only once or twice
- Has had a blood transfusion in the UK before September 1991 or received any blood products before 1986
- Has received medical or dental treatment in countries where infection control may be poor
- Is the child of a mother with HCV
- Is a regular sexual partner of someone with HCV
- Has been accidentally exposed to blood where there is a risk of transmission of HCV
- Has had tattoos, piercings, acupuncture or electrolysis where infection control procedures are poor

How do I test for HCV?

The primary screening test is a blood test for antibodies to the virus (anti-HCV), which indicates if a person has ever been infected with HCV. A positive test should be confirmed by testing a second sample. It can take three months for antibodies to become detectable. A negative test should be repeated if the exposure was within three months of the test.

About 20–40% of people will clear the virus naturally, so a test to detect HCV RNA is required to establish if the patient is still infected.

Pre-test discussion

Pre-test discussion should include:

- Hepatitis C, its natural history and the benefits offered by treatment
- What the test involves, testing timescale and confidentiality of results
- Assessment of exposure risks and establishing when the last risk activity took place
- Implications of a positive result for the individual and his/her family or close contacts
- What personal support network the individual may have; information about national/local organisations that provide support

It may also offer the opportunity to advise injecting drug users about harm minimisation and to offer them the hepatitis A and hepatitis B vaccine.

Post-test discussion

Post-test discussion should also include:

Negative antibody result

- Further testing will be required if the last exposure risk occurred in the preceding three month 'window period'
- Ways of avoiding infection in the future

Positive antibody result

- Positive antibody results should be followed by tests for HCV RNA. A second blood sample should be taken for confirmation
- Advise not to donate blood or carry an organ donor card

Positive HCV RNA result

- Patients should be referred to a specialist for further assessment
- Stop or reduce alcohol consumption (associated with more rapid progression of liver disease)
- Ways of avoiding infecting others
- Consider the need to test other family members or close contacts

Negative HCV RNA result

- A positive antibody and negative HCV RNA test indicates a previously resolved infection, but not immunity to further infection
- Patients who are antibody positive but HCV RNA negative should have a second HCV RNA test after 4–6 weeks to confirm their negative status

Hepatitis **C** Get tested. Get treated.

Hepatitis C

What you need to know


Treatment for HCV

The National Institute for Health and Clinical Excellence (NICE) recommends treatment of chronic hepatitis C with combined pegylated interferon and ribavirin, which can successfully clear the virus in up to 55% of patients overall. Current injecting drug users and people who drink excess alcohol are **not** precluded from treatment.

Since the recent NICE recommendations, liver biopsy need no longer be routine in assessing patients for treatment, though it may be advised for some patients.

Treatment may have side effects but these can be satisfactorily managed in most cases. Treatment may be contraindicated for some medical and psychiatric conditions.

From diagnosis to treatment – flow chart



Further information

- NHS hepatitis C awareness website: www.nhs.uk/hepc
- Hepatitis C Information Line: **0800 451 451**

If you require further copies of this title call 0300 123 1002 or visit www.orderline.dh.gov.uk and quote: 292688/Hepatitis C: Quick reference guide for primary care.

Hepatitis **C** Get tested. Get treated.