

SYPHILIS & SKIN DISEASE

PRIMARY SYPHILIS

- PAINLESS
- HARD EDGED
- 70% MEN, 50% WOMEN



PRIMARY SYPHILIS

- PAINLESS
- HARD EDGED
- 70% MEN, 50% WOMEN




PRIMARY SYPHILIS

- CAN ALSO OCCUR ON PARTS OTHER THAN GENITALIA EG LIPS, ORALLY ETC
- HIGHLY INFECTIOUS
- FIRM DISCRETE REGIONAL NODES



SECONDARY SYPHILIS

- GENERALISED NONTENDER LYMPHADENOPATHY
- PRIMARY CHANCRE REMAINS IN 15-25%
- INITIALLY SYMMETRICAL MACULAR, TRUNK AND PROXIMAL EXTREMITIES
- AFTER DAYS/WEEKS, PAPULAR OFTEN PALMS AND SOLES



SECONDARY SYPHILIS

- TINY PAPULAR FOLLICULAR SYPHILIS MAY DEVELOP
- IN 10% CONDYLOMA LATA, HIGHLY INFECTIOUS PAPULES IN MUCOCUTANEOUS JUNCTION



TERTIARY SYPHILIS

- GUMMAS MAY BE IDENTIFIED ON THE SKIN, IN THE MOUTH, AND IN THE UPPER RESPIRATORY TRACT. THEY APPEAR MOST COMMONLY ON THE LEG JUST BELOW THE KNEE BUT CAN OCCUR ANYWHERE



CONGENITAL SYPHILIS

- **MACULOPAPULAR DESQUAMATIVE RASH, ESPECIALLY ON PALMS AND SOLES**
- **THEN HUTCHINSONS NOTCHED TEETH, SABRE TIBIA ETC**

